


How to use Phexxi[®]

PHEXXI IS USED IMMEDIATELY **BEFORE**, OR UP TO 1 HOUR **BEFORE**, EACH ACT OF SEX.

Using Phexxi is as simple as 1-2-3


1 PREP

- Insert the plunger rod into the applicator
- Push until the plunger rod connects to the applicator
- Remove the pink cap




2 PUSH

- Gently insert the pre-filled applicator into your vagina as far as it will comfortably go
- Push the plunger rod to release the gel into the vagina



3 PREVENT

- Once the gel has been pushed into the vagina, remove the used pre-filled applicator from your vagina
- A small amount of gel may be left in the applicator but that's OK! You will still get the right dose



Go to [Phexxi.com](https://www.phexxi.com) to watch a video or read detailed instructions.



Not actual size.

INDICATION

Phexxi[®] is an on-demand method of birth control used to prevent pregnancy. Phexxi is not effective when used after sex.

IMPORTANT SAFETY INFORMATION

Rare cases (0.36%) of bladder and kidney infection have been reported. If you have a history of urinary tract problems that keep coming back, you should not use Phexxi.

Please see additional Important Safety Information on next page and full [Product Information](#) for Phexxi.

phexxi[®]
(lactic acid, citric acid, and
potassium bitartrate) Vaginal Gel
1.8%, 1%, 0.4%

IMPORTANT SAFETY INFORMATION (CONT'D)

Contact your healthcare provider if you are experiencing genitourinary side effects such as vaginal burning, itching, discharge, genital discomfort (including in male partners), yeast infection, urinary tract infection or bacterial vaginosis.

Phexxi does not protect against any sexually transmitted infections, including HIV. Avoid using Phexxi with a vaginal ring.

Avoid Phexxi if you or your sexual partner is allergic to lactic acid, citric acid, potassium bitartrate, or any of the ingredients in Phexxi. Stop using Phexxi if you develop an allergic reaction.

For more information about Phexxi, talk to your healthcare provider and see full Product Information.

Please report side effects by contacting Evofem Biosciences[®] toll-free at 1-833-EVFM BIO or contact FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.



Phexxi and the Phexxi logo are registered trademarks of Evofem Biosciences, Inc.

Trademarks, registered or otherwise, are the property of their respective owner(s).

© 2022 Evofem Biosciences, Inc. • EVFM-US-001934
April 2022