

phexxi[®]
(lactic acid, citric acid, and
potassium bitartrate) Vaginal Gel
1.8%, 1%, 0.4%

TAKE A PERSONAL APPROACH **with Phexxi[®]**

A NON-HORMONAL BIRTH CONTROL THAT CAN MEET THE INDIVIDUAL NEEDS OF WOMEN

Phexxi is a **first-in-class vaginal pH modulator (VPM)**—a bioadhesive gel that works by **maintaining an acidic vaginal pH** and **decreasing sperm motility to prevent pregnancy**, providing women with a non-hormonal, first-line contraceptive.^{1,2}

WHY MIGHT PHEXXI BE A GOOD OPTION FOR HER?

Because FDA-approved Phexxi²:

- is **non-hormonal**
- is **non-systemic**, and systemic exposure is not expected to lead to safety concerns
- **acts locally**, so it may be appropriate for breastfeeding women
- is **used in the moment**
- is **non-invasive** and **easily reversible**
- has **no contraindications** or black box warning

CONVENIENT AND EASY TO USE³

1 DOSE
HOUR
ACT

After application, **1 dose** of Phexxi:

- works immediately
- will last up to **1 hour**
- provides protection for **1 act** of vaginal intercourse



WATCH THE MOA VIDEO TO
SEE HOW PHEXXI WORKS!



INDICATIONS AND USAGE

Phexxi is indicated for the prevention of pregnancy in females of reproductive potential for use as an on-demand method of contraception.

LIMITATIONS OF USE

Phexxi is not effective for the prevention of pregnancy when administered after intercourse.

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Few cases (0.36%) of adverse reactions of cystitis, pyelonephritis and other upper urinary tract infection (UTI) have been reported in Phexxi clinical studies. Of these, one case of pyelonephritis was considered serious and required hospitalization. Avoid use of Phexxi in females with history of recurrent urinary tract infection or urinary tract abnormalities.

Please see full **Prescribing Information** and additional **Important Safety Information** on next page.


Phexxi[®] is the right choice for millions of women

phexxi[®]
(lactic acid, citric acid, and potassium bitartrate) Vaginal Gel
1.8%, 1%, 0.4%

Consider Phexxi for your patients who:

- are unhappy with their current contraceptive
- want an option that is free from hormones and won't cause hormonal side effects
- need a non-hormonal option that acts locally and may be used while breastfeeding
- are seeking contraception that is easily reversible and doesn't require a procedure
- want a woman-controlled option that's only used when needed

How to prescribe:



PATIENT: _____ DATE: _____

Phexxi vaginal gel (5-g applicator)

Sig: Administer 1 applicator intravaginally immediately before or up to 1 hour before EACH act of vaginal intercourse as needed.

Dispense: # 12 applicators (60 g total)

NDC: 69751-100-12 REFILL: **11**



IMPORTANT SAFETY INFORMATION (cont'd) ADVERSE REACTIONS

Most common adverse reactions were vulvovaginal burning sensation, pruritus, mycotic infection, genital discomfort (including male partners); urinary tract infection, bacterial vaginosis, vaginal discharge, and dysuria. Patients should consult their healthcare provider for severe or prolonged genital irritation or urinary tract symptoms.

Advise the patient:

- **To discontinue Phexxi if they develop a local hypersensitivity reaction.**
- **That Phexxi does not protect against HIV infection or other sexually transmitted infections.**
- **Avoid Phexxi use with vaginal rings.**

To report SUSPECTED ADVERSE REACTIONS, contact Evofem at toll-free phone 1-833-EVFM BIO or you may contact FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full **Prescribing Information** and additional **Important Safety Information** on previous page.

References: 1. Phexxi[®] Vaginal Gel. Medi-Span[®]; June 8, 2020. 2. Phexxi[®] [Prescribing Information]. Evofem Biosciences, Inc.: San Diego, CA; February 2022. 3. Data on file. Evofem Biosciences, Inc.